



RESEARCH ARTICLE

Article DOI: 10.21474/JNHM01/128
DOI URL: <http://dx.doi.org/10.21474/JNHM01/128>

IMPACT OF RAMADAN FASTING ON BLOOD GLUCOSE LEVELS: A SMALL PROSPECTIVE OBSERVATIONAL STUDY

Hafiz Muhammad Usman, Umna Azam, Syeda Aiman Asad and Syeda Alishba

1. Department of Physiology, Faculty of Science, University of Karachi.

Manuscript Info

Manuscript History

Received: 08 July 2025

Final Accepted: 10 August

2025 Published: September 2025

Abstract

Background:-Fasting in the month of Ramadan induces metabolic changes that may influence glucose metabolism and cardiovascular parameters in both diabetic and non-diabetic persons. This study was designed to examine the effects of Ramadan fasting on blood glucose, blood pressure, and Body Mass Index (BMI) in diabetic patients and healthy controls.

Methods:-This study was conducted at the Department of Physiology, University of Karachi from 17 February to 18 March 2026. 40 participants (20 diabetic patients and 20 healthy controls, aged 20–40 years) were included after informed consent. Demographic and lifestyle factors data were collected using a questionnaire. Fasting blood glucose, blood pressure, and BMI were measured. Fasting blood glucose was assessed after 8-10 hours of fasting using a glucometer (On Call EZ II). Data was analysed using MS Excel, and t-tests were applied with $p < 0.05$ considered significant.

Result:-In healthy controls, no significant changes were noticed in fasting blood glucose (91.5 ± 9.76 to 89.45 ± 9.85 mg/dL, $p = 0.507$), blood pressure, or BMI. In diabetic patients, fasting blood glucose significantly dropped from 169.2 ± 35.67 to 139.8 ± 33.81 mg/dL ($p = 0.008$), while changes in blood pressure and BMI were not significant.

Conclusion:-Fasting in Ramadan significantly improves glycemic control in diabetic patients without notable adverse effects on blood pressure or BMI, suggesting potential metabolic benefits under fasting conditions.

"© 2025 by the Author(s). Published by JNHM under CC BY 4.0. Unrestricted use allowed with credit to the author."

Introduction:-

Ramadan, the ninth month of Islamic lunar Hijri calendar, is observed by Muslims through fasting, known as As-Saum, which is one of the five pillars of Islam. During this time, Muslims refrain from eating, drinking, and intimate relations from dawn until sunset. Ramadan is not only a period of fasting but also a time for spiritual reflection, self-discipline, and developing empathy for those less fortunate. Ramadan fasting in is a religious obligation for Muslims, who make up about 24% of the world's 1.8 billion population (1). Globally, around 463 million adults are living with diabetes, 90% of them having type 2 diabetes (2). Pakistan has one of the highest rates of diabetes, ranking third in the world after China and India (IDF Diabetes Atlas, 10th Edition, 2021)(3) The prevalence of

Corresponding Author:-Hafiz Muhammad Usman

Address:-Department of Physiology, Faculty of Science, University of Karachi.

diabetes in Pakistan has been steadily increasing, from 11.77% in 2016 to 17.1% in 2019. In next five years, the International Diabetes Federation estimated that 26.7% of adults in Pakistan would have diabetes, expanding to approximately 33 million cases (2). Diabetes is influenced by both hereditary and lifestyle factors that include high blood pressure, high triglycerides, low HDL, cardiac diseases, diet rich in fats and carbohydrates, high alcohol intake, sedentary lifestyle, obesity, polycystic ovarian syndrome (PCOS), and older age(4). Family history also plays a serious role in the onset of diabetes. Diabetes occurs when pancreas doesn't produce enough insulin or the body can't effectively use the insulin it produces, leading to higher blood sugar levels. There are two main types of diabetes. Type 1 Diabetes (T1D), also known as insulin-dependent or juvenile diabetes, is an autoimmune condition where T-cells destroy pancreatic β cells, resulting in inadequate insulin production (5). Type 1 diabetes is not common, accounting for less than 10% of all diabetes cases, but its frequency has been growing globally since the 1950s, with an annual surge of 3-4% over the last 30 years (6). Most people with type 1 diabetes develop it before age 30, though it can occur later. Type 2 diabetes, often referred to as non-insulin-dependent or adult-onset diabetes. It's one of the leading causes of illness and death worldwide (7). Type 2 diabetes doesn't result from deficiency of insulin, but rather from cells becoming less responsive to it due to the downregulation of insulin receptors(8).

Diabetes complications can be classified into two classes: chronic complications, which develop over time, and acute complications, which can occur suddenly. A 1999-2004 data indicated prevalence of chronic microvascular complications: kidney disease, foot problems, eye damage more than macrovascular complications: heart attacks and strokes (9). Dental problems, poor immunity, pregnancy issues are other complications. Fasting has been shown to potentially reduce the risk of hypoglycaemia in type 1 diabetes, lower glycemic variability, and improve fat metabolism in both type 1 and type 2 diabetes(10). Fasting also increases the sensitivity of the body towards insulin and lowers production of glucose in the body and improves body's metabolism. Fasting, therefore, is a good approach to manage diabetes when safely accomplished under medical supervision (Herz et al., 2023). This study aimed to investigate the effect of Ramadan fasting on blood glucose levels of diabetic individuals and non-diabetic individuals.

Material and Methods:-

The study was carried out at the Department of Physiology, University of Karachi between 17 February to 18 March 2026. This study was approved from the local ethical committee of the Department. Informed consent was obtained from all participants, and they were chosen to participate in this study voluntarily. It was a small prospective observational study that included 40 participants, 20 diabetes type 2 patients and 20 non-diabetic controls, aged between 20-40 years. Demographic and lifestyle factors were recorded through a questionnaire. Healthy control participants were consuming standard normal diet. Diabetic case patients were maintaining their diet and standard medication recommended by the doctor. BMI, Blood Pressure and Blood Glucose levels were recorded at three points, start of Ramadan, mid-Ramadan, and after Ramadan. Fasting blood glucose levels were measured at 8 to 10-hours fasting period using a glucometer (On Call EZ II). It utilizes biosensor technology and designs to match standard accuracy requirements and often compared with the performance of laboratory test. Statistical analysis was conducted using MS Excel to compare findings. Continuous variables were expressed as mean \pm standard deviation and comparison was performed through T Testing and P value <0.05 was considered significant.

Results:-

This study examined the effects of fasting during Ramadan on multiple parameters, especially on blood glucose levels among control and diabetic patients and before and after the period of fasting. In healthy controls, fasting blood glucose level showed a minor decrease from 91.5 ± 9.76 mg/dL to 89.45 ± 9.85 mg/dL, and this change was not statistically significant ($p < 0.507$). Similarly, there were minimal changes in systolic and diastolic blood pressure, with 118.55 ± 11.04 to 119.4 ± 16.12 mmHg ($p < 0.827$) and 78.4 ± 12.99 to 75.9 ± 5.46 mmHg ($p < 0.422$), respectively. BMI remained stable, with no significant differences observed ($p < 0.907$). In contrast, diabetic patients experienced a notable improvement in fasting blood glucose levels, which decreased significantly from 169.2 ± 35.67 mg/dL to 139.8 ± 33.81 mg/dL ($p < 0.008$). Changes in blood pressure were less pronounced, with systolic pressure slightly reducing from 131.5 ± 15.74 to 126.1 ± 6.94 mmHg ($p = 0.314$) and diastolic pressure showing a marginal change from 82.7 ± 10.29 to 81.3 ± 4.81 mmHg ($p < 0.756$). BMI showed minor, non-significant changes ($p < 0.343$). The results are represented in Table 1.

Table 1: Comparison of Blood Glucose, Blood Pressure and BMI before and after Fasting in Diabetic and Non-Diabetic Persons

Parameters	Groups	Start of Ramadan	End of Ramadan	P- Value
Fasting Blood Glucose	Non-Diabetic	91.15±9.76	89.45±9.85	0.507
	Diabetic	169.2±35.67	139.8±33.81	0.008
Blood pressure				
Systolic	Non-Diabetic	118.55±11.04	119.4±16.12	0.827
Diastolic		78.4±12.99	75.9±5.46	0.422
Systolic	Diabetic	131.5±15.74	126.1±6.94	0.314
Diastolic		82.7±10.29	81.3±4.81	0.756
BMI (kg/m²)	Non-Diabetic	20.58±4.53	20.575±4.47	0.907
	Diabetic	26.35±2.97	26.31±5.72	0.343

Discussion:-

Ramadan month fasting, in which individuals specially Muslims abstain from food and drink (full day fasting), and few other activities from dawn until sunset, has unique effects on those living with diabetes. While fasting holds spiritual significance and potential health benefits, fasting can also pose challenges for people with diabetes. The impact of fasting on blood sugar levels and general health status depends on numerous factors, such as the type of diabetes a person has, the medications they take, their overall health, and how well they manage their condition during the month of Ramadan or fasting period. The changes in routine experienced by people with diabetes due to fasting may have potential effects on how food, insulin and other medications are processed in the body. Hence, such metabolic fluctuations may result in altered blood sugar levels that might be difficult to control without proper scheduling and monitoring. In this study, diabetic patients significantly showed lower levels of blood sugar after the period of fasting when compared with their initial levels. In non-diabetics, fasting blood glucose levels were only slightly reduced and Systolic and diastolic readings did not change much. Fasting also did not affect BMI, meaning that well monitored fasting may not change body composition.

These findings were correlatable with the study that did not find significant alterations in body weight and composition after Ramadan fasting (11). Comparable to our study, fasting in diabetic patients showed lowered fasting blood glucose, thereby suggesting glycemic control as fasting glucose was reported to decrease considerably during Ramadan (10,12). Although blood pressure showed slight reductions, these changes were not statistically significant. Weight and BMI changes were also negligible, indicating limited impact on body composition. Similar trends were noted in the study, which concluded that Ramadan fasting does not negatively affect glycemic parameters in type 2 diabetes patients and can even enhance glycemic control in some cases (13,14). Ramadan fasting offers notable glycemic benefits for diabetic individuals, particularly with appropriate medication adjustments. For non-diabetics, fasting is well-tolerated and does not disrupt metabolic constancy. However, the contrasting dietary and lifestyle patterns between these groups highlight the need for individualized and customized fasting routines and health management strategies. Furthermore, the body's ability to adapt to altered energy metabolism during Ramadan, such as increased fat oxidation, demonstrates its resilience in maintaining metabolic balance despite dietary changes (15,16). However, this study is limited by a small sample size and a short observational period that comprise of Ramadan one month duration only, which reduce the vast applicability of the outcomes.

Conclusion:-

This study explored the positive effects of Ramadan fasting on fasting blood glucose levels among healthy and especially diabetic individuals. As the Holy Prophet Muhammad (SAW) said: "Keep the fast, keep your health". Ramadan fasting may be advantageous for individuals with diabetes, particularly when shared with appropriate medication and careful diet management, hydration, and physical activity. Diabetic individuals may experience

improved blood glucose control, though the risk of blood sugar variations and dehydration remains. Characteristically, non-diabetics, showed minimal changes in studied parameters. Furthermore, healthcare experts should focus on personalized care plans for fasting during Ramadan, including steady monitoring of blood glucose. More research is required to clearly understand the long-term impacts of fasting on diabetes management and to develop much-focused guidelines for both diabetic and non-diabetic persons.

References:-

1. Bener A, Al-Hamaq AbdullaOA, Öztürk M, Çatan F, Haris P, Rajput K, et al. Effect of ramadan fasting on glycemic control and other essential variables in diabetic patients. *Ann Afr Med.* 2018;17(4):196. doi:10.4103/aam.aam_63_17
2. Azeem S, Khan U, Liaquat A. The increasing rate of diabetes in Pakistan: A silent killer. *Annals of Medicine & Surgery.* 2022 Jul;79. doi:10.1016/j.amsu.2022.103901
3. Basit A, Fawwad A, Qureshi H, Shera AS. Prevalence of diabetes, pre-diabetes and associated risk factors: second National Diabetes Survey of Pakistan (NDSP), 2016–2017. *BMJ Open.* 2018 Aug 5;8(8):e020961. doi:10.1136/bmjopen-2017-020961
4. Grundy SM, Cleeman JI, Daniels SR, Donato KA, Eckel RH, Franklin BA, et al. Diagnosis and Management of the Metabolic Syndrome. *Circulation.* 2005 Oct 25;112(17):2735–52. doi:10.1161/CIRCULATIONAHA.105.169404
5. Haller MJ, Atkinson MA, Schatz D. Type 1 Diabetes Mellitus: Etiology, Presentation, and Management. *Pediatr Clin North Am.* 2005 Dec;52(6):1553–78. doi:10.1016/j.pcl.2005.07.006
6. Patterson CC, Harjutsalo V, Rosenbauer J, Neu A, Cinek O, Skrivarhaug T, et al. Trends and cyclical variation in the incidence of childhood type 1 diabetes in 26 European centres in the 25 year period 1989–2013: a multicentre prospective registration study. *Diabetologia.* 2019 Mar 28;62(3):408–17. doi:10.1007/s00125-018-4763-3
7. Bhandari V, Dureja S, Bachhel R, Gupta M, Sidhu R. Effect of intermittent fasting on various health parameters in obese type 2 diabetics: A pilot study. *Natl J Physiol Pharm Pharmacol.* 2021;(0):1. doi:10.5455/njppp.2022.12.08281202120082021
8. Galicia-Garcia U, Benito-Vicente A, Jebari S, Larrea-Sebal A, Siddiqi H, Uribe KB, et al. Pathophysiology of Type 2 Diabetes Mellitus. *Int J Mol Sci.* 2020 Aug 30;21(17):6275. doi:10.3390/ijms21176275
9. Deshpande AD, Harris-Hayes M, Schootman M. Epidemiology of Diabetes and Diabetes-Related Complications. *Phys Ther.* 2008 Nov 1;88(11):1254–64. doi:10.2522/ptj.20080020
10. Al Hayek A, Al Zahrani WM, Al Dawish MA. Glucometric parameter changes in patients with type 2 diabetes during ramadan fasting: A prospective comparative real-world study. *Metabol Open.* 2024 Sep;23:100304. doi:10.1016/j.metop.2024.100304
11. Lessan N, Saadane I, Alkaf B, Hambly C, Buckley AJ, Finer N, et al. The effects of Ramadan fasting on activity and energy expenditure. *Am J Clin Nutr.* 2018 Jan;107(1):54–61. doi:10.1093/ajcn/nqx016
12. Elmajnoun HK, Faris ME, Abdelrahim DN, Haris PI, Abu-Median AB. Effects of Ramadan Fasting on Glycaemic Control Among Patients with Type 2 Diabetes: Systematic Review and Meta-analysis of Observational Studies. *Diabetes Therapy.* 2023 Mar 2;14(3):479–96. doi:10.1007/s13300-022-01363-4
13. Salti I, Bénard E, Detournay B, Bianchi-Biscay M, Le Brigand C, Voinet C, et al. A Population-Based Study of Diabetes and Its Characteristics During the Fasting Month of Ramadan in 13 Countries. *Diabetes Care.* 2004 Oct 1;27(10):2306–11. doi:10.2337/diacare.27.10.2306
14. Jarrar Y, Abdul-Wahab G, Mosleh R, Abudahab S, Jarrar Q, Hamdan A, et al. Does Ramadan Intermittent Fasting Affect the Fasting Blood Glucose Level among Type II Diabetic Patients? *J Clin Med.* 2023 Oct 18;12(20):6604. doi:10.3390/jcm12206604
15. el Ati J, Beji C, Danguir J. Increased fat oxidation during Ramadan fasting in healthy women: an adaptive mechanism for body-weight maintenance. *Am J Clin Nutr.* 1995 Aug;62(2):302–7. doi:10.1093/ajcn/62.2.302
16. Rahman S. Ramadan Fasting and its Health Benefits: What's New? *Open Access Maced J Med Sci.* 2022 Jul 27;10(E):1329–42. doi:10.3889/oamjms.2022.9508